

University of Groningen

New method for calculating transport coefficients and application to the Heisenberg chain

Raedt, Hans De; Raedt, Bart De

Published in:
Physical Review B

DOI:
[10.1103/PhysRevB.16.5129.2](https://doi.org/10.1103/PhysRevB.16.5129.2)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1977

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Raedt, H. D., & Raedt, B. D. (1977). New method for calculating transport coefficients and application to the Heisenberg chain. *Physical Review B*, 15(11). <https://doi.org/10.1103/PhysRevB.16.5129.2>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Errata

Erratum: Specific heat of bcc ^3He [Phys. Rev. B **15**, 2604 (1977)]

D. S. Greywall

Due to a computational error, all of the specific-heat results presented in this paper must be adjusted slightly. The correct values of the specific heat can be determined by multiplying each of the reported values by the factor $F(V)$ given in the following table. Corrected values for Θ_0 and $|J|/k_B$ are also given.

V (cm^3)	F	Θ_0 (K)	$ J /k_B$ (mK)
24.454	1.019	18.83	0.8839
23.785	1.034	20.30	0.5527
23.081	1.004	21.87	0.360
22.425	1.019	23.39	...
21.459	1.022	25.85	...

Erratum: New method for calculating transport coefficients and application to the Heisenberg chain [Phys. Rev. B **15**, 5379 (1977)]

Hans De Raedt and Bart De Raedt

Equation (4.5) should read

$$\tilde{\Omega}_q = \begin{pmatrix} 0 & (LS, LS)_q \\ (LS, LS)_q & 0 \end{pmatrix} \tilde{\chi}_q^{-1} = \begin{pmatrix} 0 & 1 \\ \frac{(LS, LS)_q}{(S, S)_q} & 0 \end{pmatrix}.$$

In Fig. 1, the temperature in the upper drawing should be $T = 40$ K.

Erratum: Crystal structure and magnetic susceptibility of $(\text{CH}_3)_2\text{NH}_2\text{MnCl}_3$ (DMMC): A low-symmetry analog of $(\text{CH}_3)_4\text{NMnCl}_3$ (TMMC) [Phys. Rev. B **13**, 3956 (1976)]

Ruth E. Caputo and Roger D. Willett

The interchain coupling J'/k should be 0.05°K , rather than 0.5°K .